

VALORES CRITICOS SELECCIONADOS DE F PARA EL ESTADÍSTICO Di DE COOK

ALFA DE 0.50													
Numerador gl= k+1													
Denominador gl = n - k- 1		2	3	4	5	6	7	8	9	10	12	15	20
	10	0.743	0.845	0.899	0.932	0.954	0.971	0.983	0.992	1	1.01	1.02	1.03
	11	0.739	0.84	0.893	0.926	0.948	0.964	0.977	0.986	0.994	1.01	1.02	1.03
	12	0.735	0.835	0.888	0.921	0.943	0.959	0.972	0.981	0.989	1	1.01	1.02
	15	0.726	0.826	0.878	0.911	0.933	0.949	0.96	0.97	0.977	0.989	1	1.01
	20	0.718	0.816	0.868	0.9	0.922	0.938	0.95	0.959	0.966	0.977	0.989	1
	24	0.714	0.812	0.863	0.895	0.917	0.932	0.944	0.953	0.961	0.972	0.983	0.994
	30	0.709	0.807	0.858	0.89	0.912	0.927	0.939	0.948	0.955	0.966	0.978	0.989
	40	0.705	0.802	0.854	0.885	0.907	0.922	0.934	0.943	0.95	0.961	0.972	0.983
	60	0.701	0.798	0.849	0.88	0.901	0.917	0.928	0.937	0.945	0.956	0.967	0.978
	120	0.697	0.793	0.844	0.875	0.896	0.912	0.923	0.932	0.939	0.95	0.961	0.972
	INFINITO	0.693	0.789	0.839	0.87	0.891	0.907	0.918	0.927	0.934	0.945	0.956	0.967

Fuente: Adaptado con permiso de Biometrika Tables for Statsticians, Vol. 1 3a. Ed., por E.S. Pearson y H.O. Hartley,eds, 1966, con autorizaci3n de losfideicomisarios de Biometrika

Pagina capturada por: M. En Adm3n. Javier Bech Vertti. Dpto. de Estadística. Universidad Aut3noma de Aguascalientes.